Welcome to Kingswood Medical Group- we work with many organisations across Swindon



Kingswood Medical Group























Eclipse deprescribing module for PIS – Reducing patient harm in moderate to severe frailty



Systolic < 115mmHg and on Hypertensive Medication (no HF or left ventricular systolic dysfunction), aged >=65 and eFI >=0.25

- Male 85yrs
- Admitted with a fall in Aug 23
- BP 100/58, 126/58 consistently low BP for the last 2-3 yrs
 - nSTEMI, CVA, recent decline, becoming more frail
 - No changes to BP meds
 - Bisoprolol, lisinopril 15mg, lercanidipine 20
- SMURF red alerts (clopidogrel and anaemia), opioid for pain, raised ACB score
 - Recent falls

Patients in this cohort have an indicative associated increased emergency admission risk of 50%.

This risk is calculated using retrospective national data for patients aged >=65 with an eFI score of >= 0.25, comparing patients in this cohort with a baseline group.



Diabetes - HbA1c < 48 and on Sulphonylurea, aged >=65 and eFI >=0.25

- Male age 91 yrs
- Hba1c = 43 (was 58 Feb 22)
- Recent falls, and deterioration, lost weight
 - Ongoing investigations
 - 120mg gliclazide a day, 2g metformin
 - BP 130/70, lisinopril was reduced 2019
 - Falls!
- Patients in this cohort have an indicative associated increased emergency admission risk of 40%.

This risk is calculated using retrospective national data for patients aged >=65 with an eFI score of >= 0.25, comparing patients in this cohort with a baseline group.



Initial thoughts

- How to contact / invite for review / needs to be done fairly quickly / followed up / will need several appts
- SMR with one of the pharmacist team
- Mark as reviewed on initial screening?
- Copy and paste evidence piece into notes, in case pt is seen by another clinician
- Falls between annual reviews/ how to ensure all clinicians aware / alert when doing blood results?
- Education

E.g. Pt appears to have a Systolic < 115mmHg and on Hypertensive Medication (no HF or left ventricular systolic dysfunction), aged >=65 and eFl >=0.25

Patients in this cohort have an indicative associated increased emergency admission risk of 50%.

This risk is calculated using retrospective national data for patients aged >=65 with an eFI score of >= 0.25, comparing patients in this cohort with a baseline group.