

Classification: Official

Publication approval reference: PAR1621_xvi

National shared care protocol:

Dronedarone for patients in adult services

Version 1

The content of the national shared care protocol was correct as of January 2022, but it has been checked against the SPC and was updated in April 25, including new pregnancy recommendations. As well these protocols, please ensure that [summaries of product characteristics](#) (SPCs), [British national formulary](#) (BNF) or the [Medicines and Healthcare products Regulatory Agency](#) (MHRA) or [NICE](#) websites are reviewed for up-to-date information on any medicine.

Specialist responsibilities

- Assess the patient and provide diagnosis; ensure that this diagnosis is within scope of this shared care protocol ([section 2](#)) and communicated to primary care.
- Use a shared decision making approach; discuss the benefits and risks of the treatment with the patient and/or their carer and provide the appropriate counselling (see [section 11](#)) to enable the patient to reach an informed decision. Obtain and document patient consent. Provide an appropriate patient information leaflet.
- Assess for contraindications and cautions (see [section 4](#)) and interactions (see [section 7](#)).
- Conduct required baseline investigations and initial monitoring (see [section 8](#)).
- Initiate and optimise treatment as outlined in [section 5](#). Prescribe the maintenance treatment for at least 4 weeks and until optimised.
- Once treatment is optimised, send the shared care agreement (or link to it) to patient's GP practice detailing the diagnosis, current and ongoing dose, any relevant test results and when the next monitoring is required. Include contact information ([section 13](#)).
- Prescribe sufficient medication to enable transfer to primary care, including where there are unforeseen delays to transfer of care.
- Conduct the required reviews and monitoring in [section 8](#) and communicate the results to primary care. After each review, advise primary care whether treatment should be continued, confirm the ongoing dose, and whether the ongoing monitoring outlined in [section 9](#) remains appropriate.

- Reassume prescribing responsibilities if a woman becomes or wishes to become pregnant.
- Provide advice to primary care on the management of adverse effects if required.

Primary care responsibilities

- Respond to the request from the specialist for shared care in writing if you do NOT wish to participate in the shared care arrangement (e.g., clinical competency reasons). It is asked that this be undertaken within 14 days of the request being made, where possible.
- If accepted, prescribe ongoing treatment as detailed in the specialists request and as per [section 5](#), taking into any account potential drug interactions in [section 7](#).
- Adjust the dose of dronedarone prescribed as advised by the specialist.
- Conduct the required monitoring as outlined in [section 9](#). Communicate any abnormal results to the specialist.
- Manage adverse effects as detailed in [section 10](#) and discuss with specialist team when required.
- Stop dronedarone and make an urgent referral to the specialist if ECG changes, hepatotoxicity, pulmonary toxicity or renal toxicity are suspected.
- Refer the management back to the specialist if the patient becomes or plans to become pregnant.
- Stop treatment as advised by the specialist.

Patient and/or carer responsibilities

- Take dronedarone as prescribed and avoid abrupt withdrawal unless advised by the primary care prescriber or specialist.
- Attend regularly for monitoring and review appointments with primary care and specialist and keep contact details up to date with both prescribers. Be aware that medicines may be stopped if they do not attend.
- Report adverse effects to their primary care prescriber. Seek immediate medical attention if they develop any symptoms as detailed in [section 11](#).
- Report the use of any over the counter medications to their prescriber and be aware they should discuss the use of dronedarone with their pharmacist before purchasing any OTC medicines.
- Avoid grapefruit juice while taking dronedarone.
- Patients of childbearing potential should take a pregnancy test if they think they could be pregnant and inform the specialist or GP immediately if they become pregnant or wish to become pregnant.

1. Background

[Back to top](#)

Dronedarone is used in the treatment of severe cardiac rhythm disorders, as a second line option when other drugs are ineffective or contraindicated. It has potentially serious adverse effects and its use requires monitoring both clinically and via laboratory testing.

Due to the significant safety concerns, NHS England (NHSE) and NHS Improvement advises that prescribers should not initiate dronedarone in primary care for any new patients. In exceptional circumstances, if there is a clinical need for dronedarone to be prescribed, this must be initiated by a specialist and only continued under a shared care arrangement in line with NICE clinical guidance ([Atrial fibrillation: NG 196](#)). Dronedarone should be used as recommended in NICE [TA 197 Dronedarone for the treatment of non-permanent atrial fibrillation](#). Where there is an existing cohort taking dronedarone, it is recommended that these patients be reviewed to ensure that prescribing remains safe and appropriate.

2. Indications

[Back to top](#)

Licensed indication: maintenance of sinus rhythm after successful cardioversion in adult clinically stable patients with paroxysmal or persistent atrial fibrillation.

[NICE TA 197](#) recommends dronedarone as an option in patients:

- whose atrial fibrillation is not controlled by first-line therapy (usually including beta-blockers), that is, as a second-line treatment option and after alternative options have been considered **and**
- who have at least 1 of the following cardiovascular risk factors:
 - hypertension requiring drugs of at least 2 different classes
 - diabetes mellitus
 - previous transient ischaemic attack, stroke or systemic embolism
 - left atrial diameter of 50 mm or greater **or**
 - age 70 years or older and
- who do not have left ventricular systolic dysfunction **and**
- who do not have a history of, or current, heart failure

3. Locally agreed off-label use

[Back to top](#)

National scoping did not identify any additional appropriate off-label indications

4. Contraindications and cautions

[Back to top](#)

This information does not replace the Summary of Product Characteristics (SPC) and should be read in conjunction with it. Please see [BNF](#) & [SPC](#) for comprehensive information.

Contraindications:

- Known hypersensitivity to dronedarone or any of the excipients
- Second- or third-degree atrio-ventricular block, complete bundle branch block, distal block, sinus node dysfunction, atrial conduction defects, or sick sinus syndrome (except when used in conjunction with a functioning pacemaker)
- Bradycardia less than 50 beats per minute
- Permanent atrial fibrillation (AF) with an AF duration ≥ 6 months (or duration unknown), and attempts to restore sinus rhythm no longer considered by the physician
- Unstable haemodynamic conditions
- History of or current heart failure, or left ventricular systolic dysfunction
- Patients with liver or lung toxicity related to previous use of amiodarone
- Co-administration with potent cytochrome P450 3A4 (CYP3A4) inhibitors, such as ketoconazole, itraconazole, voriconazole, posaconazole, telithromycin, clarithromycin, nefazodone and ritonavir (see [section 7](#))
- Co-administration with medicinal products inducing torsades de pointes, including phenothiazines, cisapride, bepridil, tricyclic antidepressants, terfenadine and certain oral macrolides (such as erythromycin), class I and III anti-arrhythmics (see [section 7](#))
- Co-administration with dabigatran
- QTc Bazett interval greater than 500 milliseconds
- Severe hepatic or renal impairment (CrCl < 30 mL/min)

Cautions:

Dronedarone can cause serious adverse reactions; clinical monitoring for development of congestive heart failure, left ventricular systolic dysfunction, QTc prolongation, liver injury, and respiratory disease are required (see also [section 8](#) & [section 9](#)).

5. Initiation and ongoing dose regimen

[Back to top](#)

- Transfer of monitoring and prescribing to primary care is normally after the patient's dose has been optimised and with satisfactory investigation results for at least 4 weeks
- The duration of treatment & frequency of review will be determined by the specialist, based on clinical response and tolerability.
- All dose or formulation adjustments will be the responsibility of the initiating specialist unless directions have been discussed and agreed with the primary care clinician
- Termination of treatment will be the responsibility of the specialist.

Initial stabilisation and maintenance dose:

400mg twice daily, with the morning and evening meals.

The starting and initial maintenance dose must be prescribed by the initiating specialist. Treatment should be initiated and monitored only under specialist supervision.

6. Pharmaceutical aspects

[Back to top](#)

Route of administration:	Oral
Formulation:	400 mg film-coated tablets
Administration details:	Tablets should be swallowed whole with a drink of water during a meal. The tablet cannot be divided into equal doses and should not be split. If a dose is missed, patients should take the next dose at the regular scheduled time and should not double the dose.
Other important information:	Grapefruit juice should be avoided during treatment with dronedarone (see section 7).

7. Significant medicine interactions

[Back to top](#)

The following list is not exhaustive. Please see [BNF](#) or [SPC](#) for comprehensive information and recommended management.

Dronedarone is associated with a large number of interactions, some of which are significant enough to contradict concurrent use, require dose adjustment and/or additional monitoring.

Dronedarone is contraindicated when co-administered with potent cytochrome P450 3A4 (CYP3A4) inhibitors, medicinal products inducing torsades de pointes, and dabigatran (see [section 4](#)).

Dronedarone is an enzyme inhibitor and can increase exposure to a number of medicines including:

- P-glycoprotein (PgP) substrates (e.g. digoxin, dabigatran, apixaban, rivaroxaban, edoxaban).
- CYP3A4 substrates (e.g. ciclosporin, statins, fentanyl, sildenafil, tacrolimus, sirolimus, everolimus, apixaban, rivaroxaban, edoxaban).
- CYP2D6 substrates (e.g. metoprolol).

Dronedarone interacts with other medicines that:

- Induce Torsade de Points or prolong qtc (e.g. Phenothiazines, cisapride, bepridil, tricyclic antidepressants, certain oral macrolides (such as clarithromycin and erythromycin), terfenadine and Class I and III anti-arrhythmics). Concomitant use is contraindicated.
- Lower heart rate (e.g. Beta-blockers, calcium channel blockers).
- Induce hypokalaemia (e.g. Diuretics, stimulant laxatives).
- Induce hypomagnesaemia (e.g. Diuretics).

Other interactions include:

- CYP3A4 inhibitors – may increase exposure to dronedarone (e.g. ketoconazole, itraconazole, voriconazole, posaconazole, ritonavir, clarithromycin, grapefruit juice). Concomitant use is contraindicated.
- Potent CYP3A4 inducers – may reduce exposure to dronedarone and are not recommended (e.g. rifampicin, phenobarbital, carbamazepine, phenytoin, St John's Wort).
- Anticoagulants – vitamin K antagonist and direct oral anticoagulant (DOAC) exposure may be increased by dronedarone (e.g. warfarin, rivaroxaban, edoxaban).

8. Baseline investigations, initial monitoring and ongoing monitoring to be undertaken by specialist

Monitoring at baseline and during initiation is the responsibility of the specialist; only once the patient is optimised on the chosen medication with no anticipated further changes expected in immediate future will prescribing and monitoring be transferred to primary care. [Back to top](#)

Baseline investigations:

- Liver function tests (LFTs)
- Urea and electrolytes (U&Es), including potassium, magnesium, and serum creatinine
- Electrocardiogram (ECG)

Initial monitoring:

- Liver function tests: after 7 days of treatment, after 1 month of treatment, then monthly until prescribing is transferred to primary care
- Urea and electrolytes: after 7 days of treatment, and after a further 7 days if any elevation is observed. If serum creatinine continues to rise then consideration should be given to further investigation and discontinuing treatment.
- Monitor concurrent medicines as appropriate, e.g. anticoagulants, digoxin.

Ongoing monitoring:

- ECG, at least every six months
- Chest X-ray and pulmonary function tests, if respiratory symptoms or toxicity suspected
- After each review, advise primary care whether treatment should be continued, confirm the ongoing dose, and whether the ongoing monitoring outlined in [section 9](#) remains appropriate.

9. Ongoing monitoring requirements to be undertaken by primary care

[Back to top](#)

See [section 10](#) for further guidance on management of adverse effects/responding to monitoring results. Monitoring information for healthcare professionals:

<https://www.sps.nhs.uk/medicines/dronedarone/>

Monitoring	Frequency
Urea and electrolytes (including magnesium and potassium) and creatinine clearance.	Every 6 months
Liver function tests	<ul style="list-style-type: none"> • Monthly for the first 6 months of treatment, and at month 9 and month 12 • Every 6 months thereafter
Symptoms of heart failure, e.g. development or worsening of weight gain, dependent oedema, or dyspnoea	Ongoing
ECG (monitoring may be conducted in primary care where this service is available)	At least every six months

(If relevant) If monitoring results are forwarded to the specialist team, please include clear clinical information on the reason for sending, to inform action to be taken by secondary care.

10. Adverse effects and other management

[Back to top](#)

Any serious adverse reactions should be reported to the MHRA via the Yellow Card scheme. Visit <https://yellowcard.mhra.gov.uk/>

For information on incidence of ADRs see relevant summaries of product characteristics.

Result	Action for primary care
As well as responding to absolute values in laboratory tests, a rapid change or a consistent trend in any value should prompt caution and extra vigilance	
Renal function: Electrolyte deficiency: hypokalaemia / hypomagnesaemia	Continue dronedarone. Correct deficiency as per local guidelines.
Creatinine elevated from baseline	Stop dronedarone for any elevations of serum creatinine which occur after transfer to primary care. Discuss urgently with specialist
Creatinine clearance <30 mL/minute/ 1.73m ²	Stop dronedarone and refer urgently to the specialist.
Cardiovascular: Bradycardia: <ul style="list-style-type: none"> Heart rate 50 - 60bpm without symptoms 	Continue dronedarone. Repeat monitoring. No action required if heart rate remains >50 without symptoms.
Heart rate ≤ 50bpm or ≤ 60bpm with symptoms	Discuss with specialist team; dose reduction may be required.
Worsening of arrhythmia, new arrhythmia, or heart block	Stop dronedarone. Urgent referral to specialist team.
Recurrence of atrial fibrillation	Refer to specialist team; discontinuation should be considered. Discontinue dronedarone if patient develops permanent AF with a duration of six months or more.
Signs or symptoms of congestive heart failure, e.g. weight gain, dependent oedema, or increased dyspnoea.	Stop dronedarone if congestive heart failure is suspected and refer urgently to specialist team.
Hepatotoxicity:	

Serum transaminases >5xULN or any symptoms of hepatic injury	Stop dronedarone. Urgent referral to initiating specialist and hepatologist.
ALT elevated >3xULN but no symptoms of hepatic injury	Continue dronedarone and repeat LFTs in 48-72 hours. If still elevated stop dronedarone and discuss with specialist urgently.
Symptoms of hepatic injury (e.g. hepatomegaly, weakness, ascites, jaundice)	Check LFTs urgently; proceed as above.
Pulmonary toxicity: new/worsening cough, shortness of breath or deterioration in general health (e.g. fatigue, weight loss, fever)	Continue dronedarone. Urgent referral to initiating specialist and respiratory specialist.
Gastrointestinal disturbance: diarrhoea, nausea, vomiting, abdominal pain, dyspepsia	Continue dronedarone. May require dose reduction; discuss with specialist if persistent.
General disorders: fatigue, asthenia	Continue dronedarone. May require dose reduction; discuss with specialist.
Dermatological disorders: rashes, pruritus, photosensitivity	Continue dronedarone. Reinforce appropriate self-care, including sun avoidance and purchasing of a broad-spectrum sunscreen (at least SPF30) if photosensitivity occurs. May require dose reduction; discuss with specialist.

11. Advice to patients and carers

[Back to top](#)

The specialist will counsel the patient with regard to the benefits and risks of treatment and will provide the patient with any relevant information and advice, including patient information leaflets on individual medicines.

The patient should be advised to report any of the following signs or symptoms to their primary care prescriber without delay:

- **Signs or symptoms of pulmonary toxicity**, e.g. breathlessness, non-productive cough or deterioration in general health (e.g. fatigue, weight loss, fever)
- **Signs or symptoms of liver injury**, e.g. abdominal pain, loss of appetite, nausea, vomiting, fever, malaise, fatigue, itching, dark urine, or yellowing of skin or eyes
- **Signs or symptoms of heart failure**, e.g. development or worsening of weight gain, dependent oedema, or dyspnoea

- **Signs or symptoms of bradycardia**, e.g. dizziness, fatigue, fainting, shortness of breath, chest pain or palpitations, confusion or trouble concentrating

The patient should be advised:

- Avoid grapefruit and grapefruit juice while taking dronedarone.
- If taking a statin and dronedarone, to report any signs of unexplained muscle pain, tenderness, weakness or dark coloured urine.
- Photosensitivity is an uncommon side effect of dronedarone (less than 1 in 100 people). If it occurs, patients should be advised on appropriate self-care: e.g. sun avoidance, protective clothing, avoiding tanning (including tanning beds) and to purchase and use of a wide broad-spectrum sunscreen (at least SPF30). These measures should be continued for the duration of therapy.

Patient information:

British Heart Foundation – Anti-arrhythmics:

<https://www.bhf.org.uk/informationsupport/heart-matters-magazine/medical/drug-cabinet/anti-arrhythmics>

12. Pregnancy, paternal exposure and breast feeding

[Back to top](#)

It is the responsibility of the specialist to provide advice on the need for contraception to male and female patients on initiation and at each review, but the ongoing responsibility for providing this advice rests with both the primary care prescriber and the specialist.

Pregnancy:

There are limited data on the use of dronedarone in pregnant women. Studies in animals have shown reproductive toxicity. Use is not recommended during pregnancy and in women of childbearing potential not using contraception.

Women of childbearing potential should use effective methods of contraception during treatment with Dronedarone and for 7 days after the final dose.

Prior to initiating Dronedarone, the prescriber should confirm that women of childbearing potential are not pregnant.

Breastfeeding:

It is unknown whether dronedarone and its metabolites are excreted in human milk. Available pharmacodynamic/toxicological data in animals have shown excretion of dronedarone and its metabolites in milk. A risk to the newborns/infants cannot be excluded. Women should be advised not to breastfeed during treatment with Dronedarone and for 7 days (about 5 half-lives) after the final dose.

A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from dronedarone therapy taking into account the benefit of breast-feeding for the child and the benefit of therapy for the woman.

13. Specialist contact information

[Back to top](#)

RUH/GWH/SFT: Via Cinapsis

14. Additional information

[Back to top](#)

Where patient care is transferred from one specialist service or GP practice to another, a new shared care agreement must be completed. Ensure that the specialist is informed in writing of any changes to the patient's GP or their contact details.

15. References

[Back to top](#)

- eBNF accessed via www.medicinescomplete.com on 12/04/2021
- Dronedarone hydrochloride 400 mg film-coated tablets (Multaq®). Sanofi. Date of revision of the text: 02/06/2020. Accessed via <https://www.medicines.org.uk/emc/product/497/> on 15/04/2025.
- Dronedarone hydrochloride 400 mg film-coated tablets (Dronedarone Aristo). Aristo Pharma. Date of revision of the text: 14/10/2020. Accessed via [Dronedarone Aristo 400mg film-coated tablets - Summary of Product Characteristics \(SmPC\) - \(emc\) | 10924](#) on 15/04/2025
- NHS England and NHS Clinical Commissioners. Aug 2019. <https://www.england.nhs.uk/publication/items-which-should-not-be-routinely-prescribed-in-primary-care-guidance-for-ccgs/> Accessed 09/04/2020
- MHRA. Drug Safety Update volume 5 issue 3: A1. October 2011. Dronedarone (Multaq ▼): cardiovascular, hepatic and pulmonary adverse events – new restrictions and monitoring requirements. Accessed via <https://www.gov.uk/drug-safety-update/dronedarone-multaq-cardiovascular-hepatic-and-pulmonary-adverse-events-new-restrictions-and-monitoring-requirements> on 09/04/2021
- NICE. TA197: Dronedarone for the treatment of non-permanent atrial fibrillation. Last updated December 2012. Accessed via <https://www.nice.org.uk/guidance/ta197> on 12/04/2021.
- NICE. NG196: Atrial fibrillation: diagnosis and management. Last updated June 2021. Accessed via <https://www.nice.org.uk/guidance/ng196> on 28/04/21.
- Specialist Pharmacy Service- Medicines Monitoring. Published July 2021 Accessed via [Dronedarone monitoring – SPS - Specialist Pharmacy Service – The first stop for professional medicines advice](#) on 24/06/2022.

- Specialist Pharmacy Service. Lactation Safety Information: dronedarone. Last reviewed 05/08/20. Accessed via <https://www.sps.nhs.uk/medicines/dronedarone/> on 12/04/2021.
- LiverTox. Dronedarone. Last updated 05/01/2018. Accessed via <https://www.ncbi.nlm.nih.gov/books/NBK548208/> 12/04/2021.
- CredibleMeds. QTDrugs List. Clarithromycin. Last updated 31st March 2021. Accessed via <https://crediblemeds.org/> on 26/04/21

16. Other relevant national guidance

[Back to top](#)

- Shared Care for Medicines Guidance – A Standard Approach (RMOC). Available from <https://www.sps.nhs.uk/articles/rmoc-shared-care-guidance/>
- NHSE guidance – Responsibility for prescribing between primary & secondary/tertiary care. Available from <https://www.england.nhs.uk/publication/responsibility-for-prescribing-between-primary-and-secondary-tertiary-care/>
- General Medical Council. Good practice in prescribing and managing medicines and devices. Shared care. Available from <https://www.gmc-uk.org/ethical-guidance/ethical-guidance-for-doctors/good-practice-in-prescribing-and-managing-medicines-and-devices/shared-care>
- NICE NG197: Shared decision making. Last updated June 2021. <https://www.nice.org.uk/guidance/ng197/>.

APC board date: 22/5/25
Last updated: 27/5/25